



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Therefore

$$0=u_0=u_2=u_4=\dots$$

That is, if a locus symmetrical with respect to the origin contains one term of odd degree, it contains no absolute term and no terms of even degree. A further expansion of the problem leads to the theorem that if a locus symmetrical with respect to the origin contains one term of even degree, all the terms are of even degree.

## CALCULUS.

239. Proposed by L. H. MacDONALD, A. M., Ph. D., Sometime Tutor in the University of Cambridge, Jersey City, N. J.

Of all triangles inscribed in a circle, find that which has the greatest perimeter.

Solution by C. N. SCHMALL, 89 Columbia Street, New York City, and REV. J. H. MEYER, S. J., Augusta, Ga.

Let  $ABC$  be the required triangle;  $O$ , the center of the given circle;  $BK=2r$ , the diameter of the circle;  $\angle BAC=\phi$ , and  $\angle BCA=\psi$ . Then the perimeter,  $p=AB+BC+AC=\text{maximum}\dots(1)$ .

But  $AB \cdot BC=2r \cdot BD$ , whence  $BC=2r \frac{BD}{AB}=2r \sin \phi$ .

Also  $\frac{AB}{BC}=\frac{\sin \psi}{\sin \phi}$ , whence  $AB=BC \frac{\sin \psi}{\sin \phi}$ , which, from the previous equation,  $=2r \sin \psi$ . We also have, from the law of signs,

$$AC=\frac{\sin(\phi+\psi)}{\sin \phi} BC=2r \sin(\phi+\psi).$$

Substituting in (1), we have

$$p=2r[\sin \phi + \sin \psi + \sin(\phi+\psi)]\dots(2).$$


Hence, for a maximum or minimum,

$$\frac{\partial p}{\partial \phi}=\cos \phi + \cos(\phi+\psi)=0, \text{ and } \frac{\partial p}{\partial \psi}=\cos \psi + \cos(\phi+\psi)=0.$$

Hence,  $\cos \phi=\cos \psi$  and  $\phi=\psi$ . Since  $\cos \phi + \cos(\phi+\psi)=0$ , we have, by substitution,  $\cos \phi + \cos 2\phi=0$ , or  $\cos \phi + 2\cos^2 \phi - 1=0$ , whence,  $\cos \phi=\frac{1}{2}$ , and therefore  $\phi=60^\circ=\psi$ .

Hence, the triangle is equilateral.

It is easy to show that  $\frac{\partial^2 p}{\partial \phi^2} \cdot \frac{\partial^2 p}{\partial \psi^2} > \frac{\partial^2 p}{\partial \phi \partial \psi}$ , and that, therefore, the triangle is a maximum.



Also solved by A. F. Carpenter, G. B. M. Zerr, A. H. Holmes, J. Scheffer, and G. W. Greenwood.

Professors Zerr, Greenwood, and Scheffer, and Mr. Holmes denoted the angles at the center subtended by the sides, by  $2\theta$ ,  $2\phi$ , and  $2\psi$ , and showed that these angles are equal to  $120^\circ$  each. Professor Carpenter showed that the inscribed triangle with one side constant and of maximum perimeter is isosceles, and then showed that of all isosceles triangles inscribed in a circle, the equilateral has the maximum perimeter.

240. Proposed by L. MORDELL, Philadelphia, Pa.


Show that the osculating conic of the catenary  $y=c \cosh \frac{x}{c}$  at the point for which  $y=\frac{c\sqrt{10}}{2}$  is a parabola.

I. Solution by J. SCHEFFER, A. M., Kee Mar College, Hagerstown, Md.

Let  $O$  be the center of curvature of the point considered, and  $C$  the center of the conic of closed contact, then (*vide* Joseph Edwards' *Differential Calculus*, where the problem is proposed as an exercise):

$$\frac{\cos \phi}{R} = \frac{1}{\rho} - \frac{\partial \phi}{\partial s}, \text{ and } \phi = \tan^{-1} \frac{\partial \rho}{\partial s},$$

where  $\phi = \angle OPC$ ,  $OP = \rho =$  radius of curvature,  $s$  an arc of the given curve, and  $PC = R$ . From



$$y = \frac{c}{2}(e^{x/c} + e^{-(x/c)}), \text{ we find } \frac{\partial y}{\partial x} = \frac{1}{2}(e^{x/c} - e^{-(x/c)}),$$

$$\frac{\partial^2 y}{\partial x^2} = \frac{1}{2c}(e^{x/c} + e^{-(x/c)}), \quad \frac{\partial s}{\partial x} = \frac{1}{2}(e^{x/c} + e^{-(x/c)}).$$

$$\therefore \rho = \left[ \left( \frac{\partial s}{\partial x} \right)^3 / \left( \frac{\partial^2 y}{\partial x^2} \right) \right] = \frac{c}{4}(e^{x/c} + e^{-(x/c)})^2. \quad \therefore \frac{\partial \rho}{\partial s} = \frac{\partial \rho}{\partial x} \cdot \frac{\partial x}{\partial s} = e^{x/c} - e^{-(x/c)}$$

therefore,  $\phi = \tan^{-1} \frac{e^{x/c} - e^{-(x/c)}}{3}$ , or,  $e^{x/c} - e^{-(x/c)} = 3 \tan \phi$ .

$$\therefore \frac{\partial \phi}{\partial x} = \frac{(e^{x/c} + e^{-(x/c)}) \cos^2 \phi}{3c}. \quad \therefore \frac{\partial \phi}{\partial s} = \frac{\partial \phi}{\partial x} \cdot \frac{\partial x}{\partial s} = \frac{2}{3c} \cos^2 \phi.$$

For the value  $y = \frac{c}{2} \sqrt{10}$ , we find  $\tan \phi = \frac{1}{3} \sqrt{6}$ ,  $\cos^2 \phi = \frac{3}{5}$ ,  $\rho = \frac{5c}{2}$ ,  $\frac{\partial \rho}{\partial s}$

$$= \frac{2}{5c}; \text{ substituting in } \frac{\cos \phi}{R} = \frac{1}{\rho} - \frac{\partial \phi}{\partial s}, \text{ we find } \frac{1}{R} \sqrt{\frac{3}{5}} = \frac{2}{\rho c} - \frac{2}{\rho c} = 0; \quad \therefore R = \infty,$$

and since the conic, the center of which is at an infinite distance, is a parabola, the assertion is proved.